

Important Formulas

➤ Distance $S = v_{av} \times t$

➤ Average acceleration

$$\text{Average acceleration} = \frac{\text{Change in velocity}}{\text{Time taken}}$$

$$a_{av} = \frac{v_f - v_i}{t}$$

➤ First Equation of Motion

$$v_f = v_i + at$$

➤ Second Equation of Motion

$$S = v_i t + \frac{1}{2} at^2$$

➤ Third Equation of Motion

$$2aS = v_f^2 - v_i^2$$

➤ To convert ms^{-1} to kmh^{-1} multiply speed with 3.6➤ To convert kmh^{-1} to ms^{-1} multiply speed with $\frac{10}{36}$ ➤ First Equation of Motion Body Moving Under Gravity $v_f = v_i + gt$ ➤ Second Equation of Motion Body Moving Under Gravity $h = v_i t + \frac{1}{2} gt^2$ ➤ Third Equation of Motion Body Moving Under Gravity $2gh = v_f^2 - v_i^2$ ➤ For bodies falling down freely value of g is positive and $v_i = 0$ ➤ For bodies moving upward value of g is negative and $v_f = 0$


2.1 Draw the representative lines of the following vectors:

(a) A velocity of $400 ms^{-1}$ making an angle of 60° with x -axis.(b) A force of $50 N$ making an angle of 120° with x -axis.


Solution

(a) A velocity of $400 ms^{-1}$ making an angle of 60° with x -axis.(i) Draw horizontal and vertical lines to represent x -axis and y -axis as shown in figure (a).

(ii) Select a suitable scale

If $100 ms^{-1} = 1 cm$
then $400 ms^{-1} = 4 cm$ (iii) Draw $4 cm$ line OQ at angle of 60° with positive x -axis. The OQ is vector \vec{V} .(b) A force of $50 N$ making an angle of 120° with x -axis.(i) Draw horizontal and vertical lines to represent x -axis and y -axis as shown in figure (b).

(ii) Select a suitable scale

If $10 N = 1 cm$ then $50 N = 5 cm$ (iii) Draw $5 cm$ line OQ at angle of 120° with x -axis. The OQ is vector \vec{F} .2.2 A car is moving with an average speed of $72 kmh^{-1}$. How much time will it take to cover a distance of $360 km$?

Given Data

$$\text{Average speed of car} = v_{av} = 72 kmh^{-1}$$

$$\text{Distance} = S = 360 km$$

To Find

$$\text{Time} = t = ?$$

Solution

By using formula of distance

$$S = v_{av} \times t$$

$$360 = 72 \times t$$

$$\frac{360}{72} = t$$

$$5 = t$$

$$t = 5 hr$$

2.3 A truck starts from rest. It reaches a velocity of $90 kmh^{-1}$ in $50 seconds$. Find its average acceleration.

Given Data

$$\text{Initial velocity of truck} = v_i = 0 kmh^{-1}$$

$$\text{Final velocity} = v_f = 90 kmh^{-1}$$

$$v_f = 90 \times \frac{10}{36} ms^{-1}$$

$$v_f = 25 ms^{-1}$$

$$\text{Time} = t = 50 s$$

To Find

$$\text{Average acceleration} = a_{av} = ?$$

Solution

By using formula of average acceleration

$$a_{av} = \frac{v_f - v_i}{t}$$

$$a_{av} = \frac{25 - 0}{50}$$

$$a_{av} = \frac{25}{50}$$

$$a_{av} = 0.5 ms^{-2}$$

2.4 A car passes a green traffic signal while moving with a velocity of $5 ms^{-1}$. It then accelerates to $1.5 ms^{-2}$. What is the velocity of car after $5 seconds$?

Given Data

$$\text{Initial velocity of car} = v_i = 5 ms^{-1}$$

$$\text{Acceleration} = a = 1.5 ms^{-2}$$

$$\text{Time} = t = 5 s$$

To Find

$$\text{Final Velocity} = v_f = ?$$

Solution

By using first equation of motion

$$\begin{aligned}v_f &= v_i + at \\v_f &= 5 + (1.5)(5) \\v_f &= 5 + 7.5 \\v_f &= 12.5 \text{ ms}^{-1}\end{aligned}$$

2.5 A motorcycle initially travelling at 18 kmh^{-1} accelerates at constant rate of 2 ms^{-2} . How far will the motorcycle go in 10 seconds?

Given Data

$$\begin{aligned}\text{Initial velocity} &= v_i = 18 \text{ kmh}^{-1} \\v_i &= 18 \times \frac{10}{36} \text{ ms}^{-1} \\v_i &= 5 \text{ ms}^{-1} \\ \text{Acceleration} &= a = 2 \text{ ms}^{-2} \\ \text{Time} &= t = 10 \text{ s}\end{aligned}$$

To Find

$$\text{Distance moved} = S = ?$$

Solution

By using second equation of motion

$$\begin{aligned}S &= v_i t + \frac{1}{2} a t^2 \\S &= (5)(10) + \frac{1}{2} (2)(10)^2 \\S &= 50 + \frac{1}{2} (2)(100) \\S &= 50 + 100 \\S &= 150 \text{ m}\end{aligned}$$

2.6 A wagon is moving on the road with a velocity of 54 kmh^{-1} . Brakes are applied suddenly. The wagon covers a distance of 25 m before stopping. Determine the acceleration of the wagon.

Given Data

$$\begin{aligned}\text{Initial velocity of wagon} &= v_i = 54 \text{ kmh}^{-1} \\v_i &= 54 \times \frac{10}{36} \text{ ms}^{-1} \\v_i &= 15 \text{ ms}^{-1} \\ \text{Distance covered} &= S = 25 \text{ m} \\ \text{Final velocity} &= v_f = 0 \text{ ms}^{-1}\end{aligned}$$

To Find

$$\text{Acceleration} = a = ?$$

Solution

By using third equation of motion

$$\begin{aligned}2aS &= v_f^2 - v_i^2 \\2(a)(25) &= (0)^2 - (15)^2 \\50(a) &= 0 - 225 \\a &= \frac{-225}{50} \\a &= -4.5 \text{ ms}^{-2}\end{aligned}$$

2.7 A stone is dropped from a height of 45 m. How long will it take to reach the ground? What will be its velocity just before hitting the ground?

Given Data

$$\begin{aligned}\text{Height} &= h = 45 \text{ m} \\ \text{Initial velocity} &= v_i = 0 \text{ ms}^{-1} \\ \text{Acceleration due to gravity} &= g = 10 \text{ ms}^{-2} \\ \text{Time} &= t = 5 \text{ s}\end{aligned}$$

To Find

$$\begin{aligned}\text{Time to reach ground} &= t = ? \\ \text{Velocity just before hitting ground} &= v_f = ?\end{aligned}$$

Solution

By using second equation of motion body moving under gravity

$$\begin{aligned}h &= v_i t + \frac{1}{2} g t^2 \\45 &= (0)(t) + \frac{1}{2} (10)(t)^2 \\45 &= 0 + 5(t)^2 \\45 &= 5(t)^2 \\\frac{45}{5} &= t^2 \\9 &= t^2 \\\sqrt{9} &= \sqrt{t^2} \\3 &= t \\t &= 3 \text{ s}\end{aligned}$$

Now for final velocity by using first equation of motion under gravity

$$\begin{aligned}v_f &= v_i + gt \\v_f &= 0 + (10)(3) \\v_f &= 0 + 30 \\v_f &= 30 \text{ ms}^{-1}\end{aligned}$$

2.8 A car travels 10 km with an average velocity of 20 ms^{-1} . Then it travels in the same direction through a diversion at an average velocity of 4 ms^{-1} for the next 0.8 km. Determine the average velocity of the car for the total journey.

Given Data

$$\begin{aligned}\text{Distance traveled} &= S_1 = 10 \text{ km} \\S_1 &= 10 \times 10^3 \text{ m} \\S_1 &= 10000 \text{ m} \\ \text{Average velocity} &= v_1 = 20 \text{ ms}^{-1} \\ \text{Next distance traveled} &= S_2 = 0.8 \text{ km} \\S_2 &= 0.8 \times 10^3 \text{ m} \\S_2 &= 800 \text{ m} \\ \text{Average velocity} &= v_2 = 4 \text{ ms}^{-1}\end{aligned}$$

To Find

$$\text{Average velocity for total journey} = v_{av} = ?$$

Solution

For S_1 time taken by using formula $S = vt$

$$\begin{aligned}t_1 &= \frac{S_1}{v_1} \\t_1 &= \frac{10000}{20} \\t_1 &= 500 \text{ s}\end{aligned}$$

For S_2 time taken

$$\begin{aligned}t_2 &= \frac{S_2}{v_2} \\t_2 &= \frac{800}{4} \\t_2 &= 200 \text{ s}\end{aligned}$$

$$\begin{aligned}\text{Total time} &= t = t_1 + t_2 \\t &= 500 + 200 \\t &= 700 \text{ s}\end{aligned}$$

$$\begin{aligned}\text{Total distance} &= S = S_1 + S_2 \\S &= 10000 + 800\end{aligned}$$

$$S = 10800 \text{ m}$$

Now by using formula of distance

$$S = v_{av} \times t$$

$$10800 = v_{av} \times 700$$

$$\frac{10800}{700} = v_{av}$$

$$15.4 = v_{av}$$

$$v_{av} = 15.4 \text{ ms}^{-1}$$

2.9 A ball is dropped from the top of a tower. The ball reaches the ground in 5 seconds. Find the height of the tower and the velocity of the ball with which it strikes the ground.

Given Data

$$\text{Time taken} = t = 5 \text{ s}$$

$$\text{Initial velocity} = v_i = 0 \text{ ms}^{-1}$$

$$\text{Acceleration due to gravity} = g = 10 \text{ ms}^{-2}$$

To Find

$$\text{Height of tower} = h = ?$$

$$\text{Final velocity} = v_f = ?$$

Solution

By using second equation of motion body moving under gravity

$$h = v_i t + \frac{1}{2} g t^2$$

$$h = (0)(5) + \frac{1}{2} (10)(5)^2$$

$$h = 0 + (5)(25)$$

$$h = 125 \text{ m}$$

Now for final velocity by using first equation of motion under gravity

$$v_f = v_i + g t$$

$$v_f = 0 + (10)(5)$$

$$v_f = 50 \text{ ms}^{-1}$$

2.10 A cricket ball is hit so that it travels straight up in the air. An observer notes that it took 3 seconds to reach the highest point. What was the initial velocity of the ball? If the ball was hit 1 m above the ground, how high did it rise from the ground?

Given Data

$$\text{Time to reach the highest point} = t = 3 \text{ s}$$

$$\text{Final velocity} = v_f = 0 \text{ ms}^{-1}$$

$$\text{Acceleration due to gravity} = g = -10 \text{ ms}^{-2}$$

To Find

$$\text{Initial velocity} = v_i = ?$$

$$\text{Height of ball 1 m above the ground} = h_t = ?$$

Solution

For initial velocity by using first equation of motion under gravity

$$v_f = v_i + g t$$

$$0 = v_i + (-10)(3)$$

$$0 = v_i - 30$$

$$30 = v_i$$

$$v_i = 30 \text{ ms}^{-1}$$

Now by using second equation of motion body moving under gravity

$$h = v_i t + \frac{1}{2} g t^2$$

$$h = (30)(3) + \frac{1}{2} (-10)(3)^2$$

$$h = 90 + (-5)(9)$$

$$h = 90 - 45$$

$$h = 45 \text{ m}$$

$$\text{Required total height} = h_t = h_{\text{gain}} + h_{\text{initial}}$$

$$h_t = 45 \text{ m} + 1 \text{ m}$$

$$h_t = 46 \text{ m}$$